arXiv:0708.4190v1 [math.GT] 30 Aug 2007

EXTERNAL EDGE CONDITION AND GROUP COHOMOLOGIES ASSOCIATED WITH THE QUANTUM CLEBSCH-GORDAN CONDITION

HAJIME FUJITA

ABSTRACT. In this article we investigate the structure of a twisted cohomology group of the first homology of a trivalent graph with a coefficient associated with its quantum Clebsch-Gordan condition. By using this investigation we shall show the existence of the external edge cocycles defined in [3] and a finite algorithm to construct such cocycles. Such cocycles are used for a combinatorial realization of the Heisenberg action on the space of conformal blocks studied by Andersen and Masbaum [1] and Blanchet, Habegger, Masbaum and Vogel [2]. Moreover we shall give a characterization of their cohomology classes in our combinatorial setting.

1. INTRODUCTION

The space of conformal blocks or a TQFT-module, which is a fundamental object in conformal field theory over Riemann surfaces or (2+1)-dimensional topological quantum field theory, has several significant symmetries including the action of the mapping class group of the surface. Andersen and Masbaum [1] and Blanchet, Habegger, Masbaum and Vogel [2] investigated the action of a certain Heisenberg group in the $SU(2)$-case, and their considerations are the starting point of what they call *spin-refined TQFT*. They studied in holomorphic and topological settings respectively. They determined the characters, and hence the isomorphism classes of the Heisenberg action. On the other hand, it is known that the dimension of the space of conformal blocks is equal to the number of certain combinatorial objects, *admissible weights of a trivalent graph*. Here we consider the dual graph of a fixed pants decomposition of the surface. In our paper [3], we constructed a combinatorial counterpart of that Heisenberg action and obtained representation matrices of the action in terms of admissible weights of the trivalent graph. In the construction we introduced a certain twisted 1-cocycle of the first homology of the graph. Namely, in [3] we showed that if there is a twisted 1-cocycle satisfying a natural condition which we call *the external edge condition* then we can construct an isomorphic Heisenberg action studied in [1] and [2]. In [3] we treated trivalent graphs, but some ingredients in it including the notion of the external edge condition can be defined for every unitrivalent graph, not only for every trivalent graph. In this article we shall show the existence of twisted 1-cocycles satisfying the external edge condition (we call it *external edge cocycles*) and a finite algorithm to construct such cocycles for a given unitrivalent graph and a positive integer called the level. Such a generalization for unitrivalent graphs leads us to a characterization of the twisted cohomology classes defined by the external edge cocycles. We shall show

---

2000 *Mathematics Subject Classification.* Primary 05C25; Secondary 81T40, 58D19.

that such classes are characterized by a certain functoriality with respect to cutting and pasting of graphs and behavior for genus one cases.

The author's research is supported by Research Fellowship of the Japan Society for the Promotion of Science for Young Scientists.

2. Notations and definitions

Let Γ be a unitrivalent graph with $3g-3+2n$ edges $\{f_l\}$, $2g-2+n$ trivalent vertices $\{v_i\}$ and n univalent vertices $\{w_m\}$. We assume that first n edges $f_1,\cdots,f_n$ have univalent vertices $w_1,\cdots,w_n$. Note that the first homology of Γ with $\mathbb{Z}_2$-coefficient is isomorphic to $\mathbb{Z}_2^g$. Fix a positive integer k and half integers $\vec{j'}=(j'_1,\cdots,j'_n)\in\{0,\frac{1}{2},\cdots,\frac{k}{2}\}^n$. Now we recall the definition of the admissible weights of level k for $(\Gamma;\vec{j'})$.

Definition 2.1 (Admissible weights)**.** A labeling of the set of edges $\vec{j}=(j_l):\{f_l\}\to\{0,\frac{1}{2},\cdots,\frac{k}{2}\}$ is an *admissible weight of level* k for $(\Gamma;\vec{j'})$ if $\vec{j}(f_l)=j'_l$ for $l=1,\cdots n$ and the following condition, which is called the *quantum Clebsch-Gordan condition of level* k, is satisfied for each trivalent vertex v_i with three edges f_{i_1}, f_{i_2} and f_{i_3};

$$\begin{cases} j_{i_1}+j_{i_2}+j_{i_3}\in\mathbb{Z} \\ |j_{i_1}-j_{i_2}|\le j_{i_3}\le j_{i_1}+j_{i_2} \\ j_{i_1}+j_{i_2}+j_{i_3}\le k. \end{cases}$$

Here we put $j_l:=\vec{j}(f_l)$. If a trivalent vertex v_i has only two edges f_{i_1} and $f_{i_2}=f_{i_3}$ then we interpret these conditions as a corresponding condition with $j_{i_2}=j_{i_3}$. Denote the set of all admissible weights of level k for $(\Gamma;\vec{j'})$ by $QCG_k(\Gamma;\vec{j'})$. Note that if $j'_1+\cdots+j'_m\notin\mathbb{Z}$ then one has $QCG_k(\Gamma;\vec{j'})=\emptyset$.

Now we define an action of $H_1(\Gamma)$ on the finite set $QCG_k(\Gamma;\vec{j'})$ as follows.

Definition 2.2. We can define an action of $H_1(\Gamma)$ on $QCG_k(\Gamma;\vec{j'})$ by;

$$\lambda:\vec{j}\mapsto\lambda\cdot\vec{j}=(j_1,\cdots,\frac{k}{2}-j_l,\cdots,j_{3g-3}),\ \text{for all } l \text{ with } f_l \text{ lying on } \lambda.$$

Indeed one can check that this operation preserves the QCG_k-condition by direct computations. Denote by $\overline{QCG_k}(\Gamma;\vec{j'})$ the quotient set with respect to this action and by $\bar{j}$ the equivalence class of $\vec{j}$.

Example 2.3. For the graph Γ in Figure 1 and $\lambda=f_3+f_4+f_5+f_6$, the action of λ on $QCG_k(\Gamma;\vec{j'})$ is given by;

$$\lambda:(j_1,j_2,j_3,j_4,j_5,j_6,j_7,j_8,j_9,j_{10})\mapsto\left(j_1,j_2,\frac{k}{2}-j_3,\frac{k}{2}-j_4,\frac{k}{2}-j_5,\frac{k}{2}-j_6,j_7,j_8,j_9,j_{10}\right).$$



Figure 1.

We define an abelian group acted on by $H_1(\Gamma)$ as follows.

Definition 2.4. Put $A(\Gamma;\vec{j'}) := (\mathbb{C}^\times)^{QCG_k(\Gamma;\vec{j'})}$ and define the action of $H_1(\Gamma)$ on $A(\Gamma;\vec{j'})$ by interchanges of the entries; $\lambda : (c_{\vec{j}}) \mapsto (c_{\lambda\cdot\vec{j}})$. By using this abelian group we can consider the twisted cochain group $C^*(\Gamma;\vec{j'}) := C^*(H_1(\Gamma);A(\Gamma;\vec{j'}))$ and its cohomology group $H^*(\Gamma;\vec{j'}) := H^*(H_1(\Gamma);A(\Gamma;\vec{j'}))$.

3. Structure of the first cohomology

Our main interest is the first cohomology group $H^1(\Gamma;\vec{j'})$, which is the group of twisted homomorphism modulo coboundaries.

Theorem 3.1. *We have the following description of the first cohomology.*

$$H^1(\Gamma;\vec{j'}) \;\cong \bigoplus_{\bar{j}\in\overline{QCG_k}(\Gamma;\vec{j'})} H_1(\Gamma)^*_{\vec{j}},$$

where we put $H_1(\Gamma)_{\vec{j}} := \{\lambda \in H_1(\Gamma) \mid \lambda\cdot\vec{j} = \vec{j}\}$ *and* $(\cdot)^* := Hom(\ \cdot\ ,\mathbb{C}^\times)$.

To show Theorem 3.1 we first show the following lemma.

Lemma 3.2. *We have the following description of the first coboundaries.*

$B^1(\Gamma;\vec{j'}) = \{\delta = (\delta_{\vec{j}}) \in Z^1(\Gamma;\vec{j'}) \mid \delta_{\vec{j}}(\lambda) = 0 \text{ for all } \vec{j} \in QCG_k(\Gamma;\vec{j'}) \text{ and } \lambda \in H_1(\Gamma)_{\vec{j}}\}$

This lemma follows from Proposition 4.7 in [3].

Proposition 3.3 ([3], Proposition 4.7)**.** *Let G be an abelian group acting on a finite set A and R an abelian group. Then G naturally acts on R^A by interchanging the entries; $g : (r_a)_{a\in A} \mapsto (r_{g\cdot a})_{a\in A}$ $(g \in G)$. Let $\tau : G \to R^A$, $\tau(g) = (\tau_a(g))$ be a twisted 1-cocycle. Then the cohomology class of τ is equal to 0 in $H^1(G;R^A)$ if and only if $\tau_a(g) = 0$ for all $a \in A$ with $g\cdot a = a$.*

Proof of Theorem 3.1. Define a homomorphism $\varphi : Z^1(\Gamma;\vec{j'}) \to \bigoplus_{\bar{j}\in\overline{QCG_k}(\Gamma;\vec{j'})} H_1(\Gamma)^*_{\vec{j}}$ by $\varphi(\delta) : \lambda \mapsto \delta_{\vec{j}}(\lambda)$ for $\delta \in Z^1(\Gamma;\vec{j'})$ and $\lambda \in H_1(\Gamma)_{\vec{j}}$. Note that $\delta_{\vec{j}}(\cdot)$ depends only on $\bar{j} \in \overline{QCG_k}(\Gamma;\vec{j'})$ over $H_1(\Gamma)_{\vec{j}}$ because of the relation

$$\begin{aligned}\delta_{\lambda'\cdot\vec{j}}(\lambda)\delta_{\vec{j}}(\lambda') &= \delta_{\vec{j}}(\lambda+\lambda') \\ &= \delta_{\lambda\cdot\vec{j}}(\lambda')\delta_{\vec{j}}(\lambda) = \delta_{\vec{j}}(\lambda')\delta_{\vec{j}}(\lambda).\end{aligned}$$

It is clear that the kernel of φ is equal to the group of coboundaries by Lemma 3.2. We show that φ is surjective. Fix $\varepsilon = (\varepsilon_{\bar{j}}) \in \bigoplus_{\bar{j}\in\overline{QCG_k}(\Gamma;\vec{j'})} H_1(\Gamma)^*_{\vec{j}}$ and take a lift $\tilde{\varepsilon}_{\bar{j}} : H_1(\Gamma) \to \mathbb{C}^\times$ of $\varepsilon_{\bar{j}} : H_1(\Gamma)_{\vec{j}} \to \mathbb{C}^\times$ for each $\bar{j}$. Define a map $\delta : H_1(\Gamma) \to A(\Gamma;\vec{j'})$ by $\delta_{\vec{j}}(\lambda) := \tilde{\varepsilon}_{\bar{j}}(\lambda)$. Then one can check that this δ is a twisted homomorphism and $\varphi(\delta) = \varepsilon$, and hence φ is surjective. □

4. Existence of the external edge cocycles

In this section we show that there is a finite algorithm to construct twisted 1-cocycles satisfying the external edge condition. That is a kind of corollary of the proof of the surjectivity of φ in Theorem 3.1. The external edge condition was defined for trivalent graphs in [3]. But it can be defined for unitrivalent graphs, and such a generalization is essential in the next section. We first introduce λ-external (resp. internal) edges.

Definition 4.1 (λ-external/λ-internal edges)**.** Let Γ be a unitrivalent graph. For $\lambda \in H_1(\Gamma)$ an edge $f_l \in \Gamma$ is said to be a λ-*external edge* if the cycle λ on Γ does not pass through f_l and one of the vertex of f_l lies on λ and the other is not. If λ does not pass through f_l and all vertices of f_l lie on λ, then we call f_l is a λ-*internal edge*. If λ does not pass through f_l and all vertices of f_l lie on λ, then we call f_l is a λ-*internal edge*. See Figure 2 for example. For $\lambda \in H_1(\Gamma)$ we denote the set of all λ-external edges by $Ex(\lambda)$.



Figure 2.

Definition 4.2 (External edge condition)**.** Fix a positive integer k. Let $(\Gamma; \vec{j}')$ be a pair of a unitrivalent graph and n half integers. Let $\delta = (\delta_{\vec{j}}) : H_1(\Gamma) \to A(\Gamma; \vec{j}')$ be a map. We say that δ satisfies the *external edge condition* if the following condition is satisfied;

$$\delta_{\vec{j}}(\lambda) = \exp\left(\pi\sqrt{-1} \sum_{f_l \in Ex(\lambda)} j_l\right) \quad \text{for all } (\lambda, \vec{j}) \text{ with } \lambda \cdot \vec{j} = \vec{j}.$$

We call a cocycle satisfying the external edge condition an *external edge cocycle* and a cohomology class represented by a external edge cocycle an *external edge class.*

Remark 4.1. For a given k and a pair $(\Gamma; \vec{j}')$, the external edge class is unique because of Lemma 3.2 (if it does exist).

Remark 4.2. If k is an odd number, then one can check that $H_1(\Gamma)_{\vec{j}} = \{0\}$, $H^1(\Gamma; \vec{j}') = \{0\}$ and the trivial cocycle satisfies the external edge condition. So we mainly concern with the case of even k.

Theorem 4.3. *For each pair $(\Gamma; \vec{j}')$ and a positive integer k there exist external edge cocycles.*

Proof. By Theorem 3.1 it is enough to check that the map $\varepsilon = (\varepsilon_{\vec{j}}) : \bigoplus_{\vec{j}} H_1(\Gamma)_{\vec{j}} \to \mathbb{C}^*$ provided by the external edge condition;

$$\varepsilon_{\vec{j}} : \lambda \mapsto \exp\left(\pi\sqrt{-1} \sum_{f_l \in Ex(\lambda)} j_l\right) \quad (\lambda \in H_1(\Gamma)_{\vec{j}})$$

is a group homomorphism. In other words we have to check that

$$\sum_{f_l \in Ex(\lambda_1+\lambda_2)} j_l \equiv \sum_{f_{l'} \in Ex(\lambda_1)} j_{l'} + \sum_{f_{l''} \in Ex(\lambda_2)} j_{l''} \bmod 2$$

for $\lambda_1, \lambda_2 \in H_1(\Gamma)_{\vec{j}}$. Note that $\varepsilon_{\vec{j}}$ depends only on $\vec{j}$. Indeed we have that for $\lambda' \in H_1(\Gamma)$

$$\sum_{f_l \in Ex(\lambda)} \Big((\lambda' \cdot \vec{j})_l - j_l \Big) = (\text{even number}) \times \left(\frac{k}{2} - j_{l_i} \right) \equiv 0 \bmod 2$$

because there are even numbers of the intersection edges $f_{l_i} \in \lambda \cap \lambda'$ and the weights on $Ex(\lambda)$ are integers because of the QCG_k-condition. To show the required relation it is enough to consider two local situations in Figure 3.



FIGURE 3.

By direct computations one can check that cancellations occur in these local situations. We only note that if the second situation occurs then we have $k \equiv 0 \bmod 4$ by the QCG_k-condition. □

Remark 4.3. According to the proof of Theorem 3.1 we have a finite algorithm to construct external edge cocycles for a given k and a pair $(\Gamma; \vec{j}')$. But we have had no canonical construction of any external edge cocycles yet.

5. CHARACTERIZATION OF THE EXTERNAL EDGE CONDITION

In this section we give a characterization of the external edge classes in our combinatorial setting.

Definition 5.1. Fix a positive integer k. For a pair $(\Gamma; \vec{j}')$ of a unitrivalent graph and n half integers denote by $\mathbb{C}(\Gamma; \vec{j}')$ the $\mathbb{C}$-vector space generated by the finite set $QCG_k(\Gamma; \vec{j}')$ and by $|\vec{j}\rangle$ a basis corresponding to $\vec{j} \in QCG_k(\Gamma; \vec{j}')$. For a twisted 1-cocycle $\delta \in Z^1(\Gamma; \vec{j}')$ we can define a homomorphism $\rho(\delta) : H_1(\Gamma) \to GL(\mathbb{C}(\Gamma; \vec{j}'))$ by $\rho(\delta)(\lambda)|\vec{j}\rangle := \delta_{\vec{j}}(\lambda)|\lambda \cdot \vec{j}\rangle$.

Proposition 5.2. *Let $\delta_1, \delta_2 : H_1(\Gamma) \to A(\Gamma; \vec{j}')$ be two twisted 1-cocycles. Assume that their cohomology classes coincide; $[\delta_1] = [\delta_2] \in H^1(\Gamma; \vec{j}')$. Then two representations $\rho(\delta_1)$ and $\rho(\delta_2)$ of $H_1(\Gamma)$ on $\mathbb{C}(\Gamma; \vec{j}')$ are isomorphic to each other.*

Proof. Fix $c = (c_{\vec{j}}) \in C^0(\Gamma; \vec{j}') = A(\Gamma; \vec{j}')$ which cobounds $\delta_1 \delta_2^{-1}$, i.e, $\delta_1(\lambda)\delta_2(\lambda)^{-1} = (\lambda \cdot c)c^{-1}$ for all $\lambda \in H_1(\Gamma)$. Then one can check that the map $\phi_c : \mathbb{C}(\Gamma; \vec{j}') \to \mathbb{C}(\Gamma; \vec{j}')$ defined by $\phi_c|\vec{j}\rangle = c_{\vec{j}}|\vec{j}\rangle$ gives an intertwiner between $\rho(\delta_1)$ and $\rho(\delta_2)$. □

Remark 5.1. By the construction we have $\phi_{c_1c_2}=\phi_{c_1}\phi_{c_2}$ for $c_1,c_2\in C^0(\Gamma;\vec{j}')$. This implies that the representation space $(\mathbb{C}(\Gamma;\vec{j}'),\rho(\delta))$ is canonically defined for the cohomology class $[\delta]$.

Consider a decomposition of the graph Γ into Γ_1 and Γ_2. (Here Γ_1 or Γ_2 may be empty. See Remark 5.2 also.) Associated with this decomposition we have the decomposition of $QCG_k(\Gamma;\vec{j}')$;

$$QCG_k(\Gamma;\vec{j}')=\bigsqcup_{\vec{j}''}QCG_k(\Gamma_1;\vec{j}_1',\vec{j}'')\times QCG_k(\Gamma_2;\vec{j}_2',\vec{j}''),$$

where $\vec{j}_1'$ (resp. $\vec{j}_2'$) is a pair of half integers which are entries of $\vec{j}'$ on the univalent vertices on Γ_1 (resp.Γ_2). In this operation we say that $(\Gamma_1;\vec{j}_1',\vec{j}'')$ and $(\Gamma_2;\vec{j}_2',\vec{j}'')$ are a decomposition of $(\Gamma;\vec{j}')$. Now for fixed $\vec{j}\in QCG_k(\Gamma_2;\vec{j}_2',\vec{j}'')$ we have an inclusion

$$\iota_{\vec{j}}:QCG_k(\Gamma_1;\vec{j}_1',\vec{j}'')\times\{\vec{j}\}\hookrightarrow QCG_k(\Gamma;\vec{j}')$$

and the induced projection

$$\iota_{\vec{j}}^*:A(\Gamma;\vec{j}')\to A(\Gamma_1;\vec{j}_1',\vec{j}'').$$

Combining with the inclusion $H_1(\Gamma_1)\hookrightarrow H_1(\Gamma)$ these data induce the cochain map (denote it by the same letter)

$$\iota_{\vec{j}}^*:C^*(\Gamma;\vec{j}')\to C^*(\Gamma_1;\vec{j}_1',\vec{j}'').$$

Remark 5.2. In this article we mean a *decomposition of the graph* by cutting some internal edges in Γ. See Figure 4 below.



FIGURE 4.

Now we define an equivalence relation $\sim_\Gamma$ between representations induced by the cocycles.

Definition 5.3 (Equivalence under the factorization)**.** We call two twisted 1-cocycles δ_1 and δ_2 are *equivalent under the factorization* if for any decomposition of $(\Gamma;\vec{j}')$ into $(\Gamma';\vec{j}_1',\vec{j}'')$ and $(\Gamma'';\vec{j}_2',\vec{j}'')$ the induced representations $\rho(\iota_{\vec{j}}^*(\delta_1))$ and $\rho(\iota_{\vec{j}}^*(\delta_2))$ on $\mathbb{C}(\Gamma';\vec{j}_1',\vec{j}'')$ are isomorphic representations for all $\vec{j}\in QCG_k(\Gamma'';\vec{j}_2',\vec{j}'')$. We denote $\delta_1\sim_\Gamma\delta_2$ if δ_1 and δ_2 are equivalent under the factorization, and we denote by $[\delta]_\Gamma$ the equivalence class of a twisted 1-cocycle δ.

Theorem 5.4. *Let* $R(\Gamma;\vec{j}')$ *be the set of equivalence classes of twisted 1-cocycles under the equivalence relation* $\sim_\Gamma$. *The map* $\rho : R(\Gamma;\vec{j}') \to H^1(\Gamma;\vec{j}')$ *defined by* $\rho([\delta]_\Gamma) = [\delta]$ *gives an isomorphism.*

Note that $R(\Gamma;\vec{j}')$ has a structure of an abelian group induced from that of $Z^1(\Gamma;\vec{j}')$; $[\delta_1]_\Gamma \cdot [\delta_2]_\Gamma := [\delta_1\delta_2]_\Gamma$. The isomorphism given in the theorem is an isomorphism between abelian groups. By definition the map ρ is surjective (if it is well-defined). To show the well-definedness and the injectivity we first show the following special case.

Proposition 5.5. *Let* $\Gamma(n)$ *be a unitrivalent graph having* n *univalent vertices and satisfying* $H_1(\Gamma(n)) \cong \mathbb{Z}_2$ *(Figure 5). Then we have that* $\rho(\delta_1)$ *and* $\rho(\delta_2)$ *are isomorphic if and only if their cohomology classes coincide;* $[\delta_1] = [\delta_2] \in H^1(\Gamma(n);\vec{j}')$.



Figure 5.

Proof. Let λ be the generator of $H_1(\Gamma(n))$. By Lemma 3.2 it is enough to show that if $\rho(\delta_1)$ and $\rho(\delta_2)$ are isomorphic then $(\delta_1)_{\vec{j}}(\lambda) = (\delta_2)_{\vec{j}}(\lambda)$ for all pairs $(\lambda, \vec{j})$ with $\lambda \cdot \vec{j} = \vec{j}$. Note that if there is such $\vec{j}$, then it is equal to $\vec{j}_0 := (\vec{j}', \frac{k}{4}, \cdots, \frac{k}{4})$. By the assumption $\rho(\delta_1)$ and $\rho(\delta_2)$ are isomorphic, in particular we have that $(\delta_1)_{\vec{j}_0}(\lambda) = Tr(\rho(\delta_1)(\lambda)) = Tr(\rho(\delta_2)(\lambda)) = (\delta_2)_{\vec{j}_0}(\lambda)$. □

Lemma 5.6. *The map* ρ *is well-defined.*

Proof. We have to check that $\delta_1 \sim_\Gamma \delta_2$ then $[\delta_1]$ is equal to $[\delta_2]$ in $H^1(\Gamma;\vec{j}')$. By Lemma 3.2 it is enough to show that $(\delta_1)_{\vec{j}}(\lambda) = (\delta_2)_{\vec{j}}(\lambda)$ for all pairs $(\lambda, \vec{j})$ with $\lambda \cdot \vec{j} = \vec{j}$. Fix such a pair $(\lambda, \vec{j})$. Consider a decomposition of Γ into two unitrivalent graph $\Gamma(\lambda)$ and $\Gamma'(\lambda)$ as follows. For $\lambda \in H_1(\Gamma)$, we first cut λ-external and λ-internal edges of Γ. Then we define $\Gamma(\lambda)$ to be the component of this graph which contains the cycle λ and $\Gamma'(\lambda)$ to be the complementary graph.



Figure 6.

Note that $\Gamma(\lambda)$ can be written $\Gamma(n')$ for some n' (or as a disjoint union of such graphs). By the assumption induced representations $\rho(\iota^*_{\vec{j}_3}(\delta_1))$ and $\rho(\iota^*_{\vec{j}_3}(\delta_2))$ on $\mathbb{C}(\Gamma(\lambda);\vec{j}',\vec{j}'_1)$ are isomorphic representations for all $\vec{j}_3\in QCG_k(\Gamma'(\lambda);\vec{j}'_2,\vec{j}'')$. In particular by Proposition 5.5 we have that

$$(\delta_1)_{\vec{j}}(\lambda)=\left(\iota^*_{\vec{j}_3}(\delta_1)\right)_{\vec{j}_0'''}(\lambda)=\left(\iota^*_{\vec{j}_3}(\delta_2)\right)_{\vec{j}_0'''}(\lambda)=(\delta_2)_{\vec{j}}(\lambda),$$

where $\vec{j}_0'''\in QCG_k(\Gamma(\lambda);\vec{j}',\vec{j}'_1)$ satisfies $\iota_{\vec{j}_3}(\vec{j}''')=\vec{j}$ and $\lambda\cdot\vec{j}_0'''=\vec{j}_0'''$. ☐

Lemma 5.7. *The homomorphism ρ is injective.*

Proof. We show that if $\rho([\delta]_\Gamma)=[\delta]=0$ then δ is equivalent to 0 under the factorization. Take a decomposition $(\Gamma_1;\vec{j}'_1,\vec{j}'')$ and $(\Gamma_2;\vec{j}'_2,\vec{j}'')$ of $(\Gamma;\vec{j}')$ arbitrary. Then we have a homomorphism $\iota^*_{\vec{j}}:H^1(\Gamma;\vec{j}')\to H^1(\Gamma_1;\vec{j}'_1,\vec{j}'')$. By the assumption we have that $\iota^*_{\vec{j}}[\delta]=0$ and hence the induced representation $\rho(\iota^*_{\vec{j}}(\delta))$ on $\mathbb{C}(\Gamma_1;\vec{j}'_1,\vec{j}'')$ is isomorphic to $\rho(0)$ by Proposition 5.2. In particular we have $\delta\sim_\Gamma 0$. ☐

By Theorem 4.3 and Lemma 3.2, there exists the unique external edge class for each pair $(\Gamma;\vec{j}')$. Denote the class by $\delta_{\rm ext}[\Gamma;\vec{j}']$. The following follows from the definition of the external edge condition.

Proposition 5.8. *The mapping $(\Gamma;\vec{j}')\mapsto\delta_{\rm ext}[\Gamma;\vec{j}']$ is functorial with respect to the factorization, that is, for every decomposition $(\Gamma_1;\vec{j}'_1,\vec{j}'')$ and $(\Gamma_2;\vec{j}'_2,\vec{j}'')$ of $(\Gamma;\vec{j}')$ we have $\iota^*_{\vec{j}}\delta_{\rm ext}[\Gamma;\vec{j}']=\delta_{\rm ext}[\Gamma_1;\vec{j}'_1,\vec{j}'']$.*

Lemma 5.9. *A functorial mapping $(\Gamma;\vec{j}')\mapsto[\delta(\Gamma;\vec{j}')]\in H^1(\Gamma;\vec{j}')$ is characterized by values at all pairs of the form $(\Gamma(n);\vec{j}')$.*

Proof. We show that if two functorial mappings $\{[\delta(\Gamma;\vec{j}')]\}$ and $\{[\delta'(\Gamma;\vec{j}')]\}$ satisfy $[\delta(\Gamma(n);\vec{j}')]=[\delta'(\Gamma(n);\vec{j}')]$ for all $(\Gamma(n);\vec{j}')$ then we have $[\delta(\Gamma;\vec{j}')]=[\delta'(\Gamma;\vec{j}')]$ for all $(\Gamma;\vec{j}')$. Fix a pair $(\Gamma;\vec{j}')$ and a cycle $\lambda\in H_1(\Gamma)$. Then by the same argument in the proof of Lemma 5.6, one can show that $\delta_{\vec{j}}(\Gamma;\vec{j}')(\lambda)=\delta'_{\vec{j}}(\Gamma;\vec{j}')(\lambda)$ for all pairs $(\lambda,\vec{j})$ with $\lambda\cdot\vec{j}=\vec{j}$. ☐

Combining with these arguments we have a characterization of the external edge classes.

Theorem 5.10. *The external edge classes are unique classes which define a functorial mapping satisfying $\iota^*_{\vec{j}}\delta_{\rm ext}[\Gamma;\vec{j}']=\delta_{\rm ext}[\Gamma_1;\vec{j}'_1,\vec{j}'']$ for any decompositions of $(\Gamma;\vec{j}')$ with $\Gamma_1=\Gamma(n')$ for some n'.*

Remark 5.3. In the case of $\Gamma=\Gamma(n)$ there is a standard external edge cocycle $\delta_{\rm ext}(\Gamma(n);\vec{j}')=(\delta_{\rm ext}(\Gamma(n);\vec{j}')_{\vec{j}})$ defined by

$$\delta_{\rm ext}(\Gamma(n);\vec{j}')_{\vec{j}}(\lambda)=\begin{cases}\exp\pi\sqrt{-1}(j'_1+\cdots+j'_n) & (\vec{j}=\vec{j}_0)\\ 1 \quad (\vec{j}\neq\vec{j}_0)\end{cases}$$

for $\vec{j}\in QCG_k(\Gamma(n);\vec{j}')$ and the generator λ of $H_1(\Gamma(n))$.

References

[1] J. E.Andersen and G. Masbaum, *Involutions on moduli spaces and refinements of the Verlinde formula*, Math. Ann., 314 (1999), 291-326.

[2] C. Blanchet, N. Habegger, G. Masbaum and P. Vogel, *Topological quantum field theories derived from the Kauffman bracket*, Topology 34, No.4 (1995), 883-927.

[3] H. Fujita, *A combinatorial realization of the Heisenberg action on the space of conformal blocks*, arXiv:math.GT/0708.3309, submitted.

Graduate school of Mathematical Sciences, the University of Tokyo, Komaba 3-8-1, Meguro-Ku, Tokyo, 153-8914 Japan
E-mail address: `hajime@ms.u-tokyo.ac.jp`